
office







Clipboard

PubMed

Nucleotide

Search PubMed

Protein Genome Structure

PopSet Taxonomy

omim Clear

Details

8

About Entrez

Display

for

Limits

Abstract Sort

Preview/Index



Clip Add Order

Text Version

Entrez PubMed
Overview
Help | FAQ
Tutorial
New/Noteworthy
E-Utilities

PubMed Services
Journals Database
MeSH Browser
Single Citation Matcher
Batch Citation Matcher
Clinical Queries
LinkOut
Cubby

Related Resources Order Documents NLM Gateway TOXNET Consumer Health Clinical Alerts ClinicalTrials.gov PubMed Central

Privacy Policy

□ 1: Cancer Lett 2002 Feb 8;176(1):47-55

Related Articles, Links

EUSEVIERSCIENCE FULL-TEXT ARTICLE

Ectopic expression of homeodomain protein CDX2 in intestinal metaplasia and carcinomas of the stomach.

History

Bai YQ, Yamamoto H, Akiyama Y, Tanaka H, Takizawa T, Koike M, Kenji Yagi O, Saitoh K, Takeshita K, Iwai T, Yuasa Y.

Department of Surgery, Graduate School of Medicine and Dentistry, Tokyo Medical and Dental University, Tokyo, Japan.

The roles of CDX2 and CDX1 homeobox genes during gastric carcinogenesis remain poorly defined. We have studied the expression of CDX2/1 in gastric cancers and intestinal metaplasia (IM) of 69 gastric carcinoma patients by immunohistochemistry. CDX2/1 were shown to be ectopically overexpressed in IM in 41 (85%) of 48, and 47 (90%) of 52 cases, respectively. The expression of CDX2/1 was detected in 38 (55%) and 51 (74%) of the 69 gastric carcinomas, respectively. The histological type of the gastric carcinomas was independently associated with CDX2 expression, but not with that of CDX1, with higher CDX2 expression in intestinal type (differentiated type) than in diffuse type (undifferentiated type) gastric carcinomas. Our results thus suggest that CDX2 and CDX1 may play a role during IM formation and gastric carcinogenesis.

PMID: 11790453 [PubMed - indexed for MEDLINE]



Write to the Help Desk
NCBI | NLM | NIH
Department of Health & Human Services